**Evolution of Volatile Compounds and Biogenic Amines throughout the Shelf-life of
Marinated and Salted Anchovies (*Engraulis encrasicolus*)**

Alexandre Dehaut [#], Charlotte Himber [#], Véronique Mulak [§], Thierry Grard [▽], Frédéric
Krzewinski [‡], Bruno Le Fur [§], Guillaume Duflos ^{#,*}

[#] ANSES, Laboratoire de Sécurité des Aliments – Département des Produits de la Pêche et de
l'Aquaculture, Boulevard du Bassin Napoléon, 62200 Boulogne-sur-Mer, France

[§] Plate-forme d'Innovation Nouvelles Vagues, 15 - 17 Rue Magenta, 62200 Boulogne-sur-Mer,
France

[▽]Unité Biochimie des Produits Aquatiques – Université du Littoral-Côte d'Opale, Boulevard du
Bassin Napoléon, 62200 Boulogne-sur-Mer, France

[‡]Université de Lille I, Unité de Glycobiologie Structurale et Fonctionnelle – UGSF, Bâtiment
C9 - Cité Scientifique, 59650 Villeneuve d'Ascq, France

*Corresponding author: E-mail: guillaume.duflos[at]anses.fr. Phone: +33 321 992 500. Fax:
+33 321 991 725

Published in Journal of Agricultural and Food Chemistry and available online at
<https://doi.org/10.1021/jf5021736>

15 ABSTRACT

16 Producers of processed anchovies have developed hazard analysis and critical control points
17 (HACCP) to guarantee the quality of their products. Nonetheless there is a lack of objective data
18 to determine products shelf-life. The quality of a product is usually established based on its
19 safety and organoleptic properties. These parameters were assessed by monitoring the profiles
20 of volatile compounds and quantitating six biogenic amines in samples of two types of
21 processed anchovies during their shelf-life. Regarding biogenic amines, quantities were below
22 the regulatory limits throughout shelf-life, except when a temperature abuse was applied for
23 marinated samples. Moreover, this work highlights an optimum volatile profile at five and six
24 months of storage for salted and marinated anchovies, respectively. This is the result of higher
25 content of six aldehydes and nine ketones compounds, mainly coming from the lipid oxidation.

26

27 **KEYWORDS:** Anchovies, Shelf-life assessment, Volatile compounds, Biogenic amines,
28 SPME-GC/MS

29 INTRODUCTION

30 Anchovies are one of the most caught fish species in the world with an estimated 3.5 million
31 tons landed in 2010¹ and quantities consumed estimated at 369,000 tons in 2008. Anchovies can
32 be processed using a wide range of techniques, as salt ripening or marination in vinegar.
33 Consumers are more and more aware of food quality and food safety. In a way to manage such
34 issues, the European Union has encouraged the application of good manufacturing practices and
35 the development of hazard analysis and critical control points (HACCP). This approach has
36 been developed in industry with specific applications for salted and marinated anchovies.

37 After their catch, anchovies can be ripened in salt or marinated in vinegar for a specified period
38 and preserved until consumption. The shelf-life of processed anchovies has only been
39 empirically estimated and there is a real lack of objective information on how the properties of
40 products change during storage. The assessment of optimal shelf-life will ensure to consumers a
41 safe product with good quality.

42 Numerous studies have described the microbial flora present in salted and marinated anchovies.
43 *Lactobacillus* and *Micrococcus* genera have been identified as the main bacteria present in
44 processed marinated anchovies,² presence of *Enterobacteriaceae* was also demonstrated.^{3,4} In
45 salted anchovies, potential histaminogenic flora, as *Enterobacteriaceae* and *Staphylococcus*, has
46 been reported.³⁻⁵ In a study on brined anchovies,⁶ halophilic and psychrophilic flora increased
47 respectively at room temperature and in chilled conditions, for samples conserved in low NaCl
48 concentrations brines. A consensus is observed from previous authors to conclude that hygiene
49 management provides a better control of these bacteria. In addition, a study on the ingredients
50 added during the marination process showed that they have no impact on microbiological
51 quality.⁷

Histamine is a biogenic amine resulting from the enzymatic decarboxylation of histidine. This molecule is responsible for many cases of food poisoning, with various symptoms.⁸ Due to the high histidine content in their flesh, numerous fish families (Scombridae, Clupeidae or Engraulidae) show a high histamine risk. This is the major food safety risk encountered in such products. European regulation (Commission Regulation (EC) No 1441/2007 & Commission Regulation (EC) No 1019/2013) has defined a limit of 400 mg.kg⁻¹ of histamine for a single products having undergone enzymatic maturation treatment in brine.^{9,10} Other biogenic amines have been reported, especially tyramine, putrescine, cadaverine, spermine and spermidine.¹¹⁻¹³ Studies tracing biogenic amines during the storage period indicate changes in tyramine, histamine and tryptamine.^{11,13,14} Very few studies have been carried out on the storage of marinated samples and focus mainly on the marination step itself.^{2,12} Furthermore, refrigeration limits the biogenic amine content.^{6,13,14} Analysis of volatile compounds in different food matrices has been increasing over the last decade.¹⁵⁻¹⁷ The volatilome of a matrix can be analyzed by trapping volatiles on a solid phase micro-extraction (SPME) fiber, in the headspace of a vial.¹⁸ At last, volatiles can be quantitated thanks to GC/MS device. To date, most studies on the volatile compounds of processed anchovies have been carried out to assess the maturation step of salted anchovies. Thus, indicators of maturation have been defined for salted anchovies as aldehydes, alkadienals, alcohols and ketones.^{19,20} Acetaldehyde, 2-methyl-butanal and 3-methyl-propanal have been described as the most odorant compounds in ripened anchovies.²¹ Based on sensory analysis, the optimal maturation of salted anchovies has been estimated at close to 160 days.²²

As anchovies processing is well documented, the goal of this study was focused on the less studied shelf-life assessment of processed anchovies. To this end, both salt-ripened and marinated anchovies stored in different conditions, with and without temperature change, were

studied. The first half of the study was focused on volatilome monitoring throughout storage whereas the other half dealt with safety issues by quantitating biogenic amines.

MATERIALS AND METHODS

Chemicals

NaCl used for the headspace analysis was obtained from Oxoid (Dardilly, France). An *n*-paraffin mix, C₅ to C₁₅, used for the calculation of the linear retention index (LRI) was acquired from Sigma-Aldrich (St-Quentin-Fallavier, France). All pure compounds used to improve identification reliability including: ethanol, dimethyl sulfide, 2-methyl-propanal, 2,3-butanedione, butanal, 3-methyl-butanal, 2-methyl-butanal, 1-penten-3-ol, 2-ethylfuran, 1-pentanol, 2,3-hexanedione, hexanal, heptanal, benzaldehyde, 1-octen-3-ol, (E,E)-2,4-heptadienal, octanal, benzeneacetaldehyde, nonanal, (E,Z)-2,6-nonadienal and decanal were purchased from Sigma-Aldrich (St-Quentin-Fallavier, France). All chemicals used for quantitating biogenic amines were obtained from Sigma-Aldrich (St-Quentin-Fallavier, France), except perchloric acid purchased from Carlo Erba (Val-de-Reuil, France), and toluene and acetonitrile from VWR (Fontenay-sous-Bois, France).

Anchovy processing and preservation

Samples of anchovies were provided by the Association of French Fish Processing Industries (*Confédération des Industries de Traitement des Produits de la Pêche Maritimes* (CITPPM - Paris)) and held at the innovation platform *PFI-Nouvelles Vagues* (Boulogne-sur-Mer, France). Samples of anchovy fillets (*Engraulis encrasicolus*) consisted of processed, packed, salted or marinated anchovies.

For salted anchovies, samples were prepared by immersion in brine. Fish were gutted and beheaded and then transferred to vats, with alternating layers of salt and anchovies. Saturated

brine was then added. Vats were covered and pressure was applied to the lids to remove trapped air. Anchovies were ripened at room temperature for several months (a minimum of 3 months). After ripening, samples were packed in glass jars.

Marinated fish samples were produced from fresh anchovies that were cleaned, beheaded, gutted and filleted. After 24 hours freezing to manage parasite risk, anchovies were then immersed in a vinegar solution. A marination period of a few days (with a minimum of 24 hours) under refrigeration was carried out. Anchovies were then covered with oil and vacuum packed.

All storage time applied in this study were set to exceed the shelf-life advocated by the guide to good hygiene practices. Two types of temperature of storage were applied to samples: with (wR) or without rupture (w/oR). Concerning marinated samples, storage without any change of temperature consisted of incubation at 4°C. Storage with breaks in the temperature conditions consisted of one-third of the time at 4°C and the remaining two-thirds at 8°C. For the salted anchovies, storage without any change of temperature consisted of incubation at 15°C, to insure their inherent shelf stability. Rupture applied in the temperature conditions consisted of two-thirds of the time at 15°C and the remaining one-third at 22°C.

Analysis of the headspace by SPME-GC/MS

Analysis of the headspace of vials containing anchovy samples was carried out as described by Duflos *et al.*¹⁷. Briefly, 50 g of flesh were added to 100 mL of NaCl solution in ultrapure water at 300 g.kg⁻¹ then ground with a stomacher at 300 rpm for 2 min. Then, 45 g of filtrate was centrifuged at 10,000 ×g, for 10 min at 4°C. The supernatant (11 mL) was then distributed in 20 mL SPME vials and tightly sealed. SPME vials were placed at 4°C on a refrigerated tray. The vials were handled using a combiPAL autosampler (CTC Analytics, Zwingen, Switzerland). The vial was placed for 10 min at 50°C and shaken at 500 rpm. A 75 µm CAR/PDMS fiber

(Supelco, Lyon, France) was introduced into the headspace of the vial to collect volatile compounds for 40 min at 50°C under the same conditions. The fiber was then removed and placed 1 min in a Merlin Microseal injector (at 250°C) in splitless mode, of a gas chromatograph GC-2010 equipped with a QP2010 Plus mass spectrometer (Shimadzu, Kyoto, Japan). The GC was equipped with a SLB-5 MS capillary column (60 m × 0.25 mm × 0.25 µm) from Supelco (St-Quentin-Fallavier, France) and helium was used as carrier gas at a flow of 1.78 mL/min. The temperature conditions applied in the oven were as follows: 5 minutes at 35°C then increased to 100°C at a rate of 10°C/min, then increased to 280°C at a rate of 20°C/min and maintained for 5 minutes. Spectrometric analysis was performed with the following conditions: interface temperature at 280°C, ionization voltage at 70 eV, mass range from 33 to 200 m/z and a scan frequency of 500 Hz. After each injection, the fiber was reconditioned at 300°C with helium flow for 10 min. Volatile compounds were automatically integrated by GC/MS Postrun analysis (Shimadzu, Kyoto, Japan), with a given retention time, a target ion and reference ions for each volatile compound (**Table 1**). A normalization of each area (*i.e.* division of the area by the sum of areas of all compounds) was realized. Compounds were identified by comparing them with mass spectra in the NIST 08 library database and with LRI found in the literature,²³ calculated using the equation in H. van den Dool and D. J. Kratz.²⁴ The headspace analysis was performed in duplicate for all samples.

Separation and quantitation of biogenic amines

Quantitative assay of biogenic amines was carried out using the method recommended by Commission Regulation (EC) No 1441/2007 with high performance liquid chromatography (HPLC).^{25,26} Briefly, to extract biogenic amines, 5 g of anchovies were added to 100 µL of 1,3-diaminopropane used as internal standard and 10 mL of 0.2 M perchloric acid. The mix was blended with an Ultraturrax homogenizer and centrifuged at 7000 ×g for 5 min at 4°C. Then,

100 μL of supernatant was transferred in 300 μL of sodium carbonate and 400 μL of dansyl chloride, homogenized and kept in the dark at 60°C for 5 min. After derivatization, 100 μL of L-proline was added to remove excess dansyl chloride and kept in the dark at room temperature for 15 min. Then, 500 μL of toluene was added; after settling, the aqueous phase was frozen and the organic phase containing dansyl derivates was specifically recovered and evaporated under nitrogen flow for 5 min. The dry residue was dissolved in 200 μL of acetonitrile, then filtered at 0.2 μm and injected in a high-performance liquid chromatograph. Biogenic amines were separated on a Kromasil C18 reverse phase column (250 mm x 4.6 mm x 5 μm) from Alltech (Deerfield, IL, USA) with a water/acetonitrile gradient and a flow of 1 mL/min. After 30 min of separation, the chromatogram showed the peaks of the six biogenic amines: putrescine, cadaverine, histamine, tyramine, spermidine and spermine and the internal standard (1,3-diaminopropane). Quantitation of biogenic amines was performed calculating each response factor against 1,3-diaminopropane and using a calibration curve.

Experimental design and statistical analysis

For the analysis of volatile compounds, marinated samples were studied for 9 months and salted samples for 18 months, to reach or exceed the shelf life recommended by good manufacturing practices. Samples were irregularly supplied by producers but a minimum of two samples was analyzed in duplicate each month for the duration of the tested shelf-life. A principal component analysis (PCA) was carried out on the data using open-source data-mining software Tanagra 1.4.²⁷ For each PCA, the two first PC were retained for interpretations. Correlations with principal components resulting from analysis determined which volatile compounds best characterized each group, as visualized by PCA.

Experiments on biogenic amines were designed as follows: for each pair of parameters (methods of preparation and storage conditions), five samples were taken for initial

measurements (M0) and three samples for remaining month. Samples were analyzed every two months for 8 months (marinated samples) and 14 months (salted anchovies). For each analysis, measures were performed in duplicate. Comparisons of average results of assays, for biogenic amines, were carried out using non-parametric Kruskal-Wallis tests and *post hoc* pairwise Wilcoxon rank sum tests because the results did not follow a normal distribution and there was not homoscedasticity of variances. The tests were performed using the package “stats” from the R software version 3.0.0. Kruskal-Wallis tests leading to significant results ($p\text{-value} < 0.05$) indicated that significant differences were observed for the concentration of the biogenic amines with respect to storage time. Pairwise Wilcoxon rank sum tests with $p\text{-values} < 0.05$ indicated that the two compared means were significantly different.

RESULTS

Selection of the volatile compounds for PCA analysis

From the whole dataset obtained by chromatogram integration, 36 compounds were selected as useful for our analysis (**Table 1**). These mostly included aldehydes, ketones and alcohols and three compounds that do not belong to these chemical classes: 2-ethylfuran (furan), dimethyl sulfide (organosulphur) and trimethylamine (TMA) (amine).

Among the compounds selected for statistical analysis, none allowed to differentiate storage conditions with or without rupture (data not shown). This study thus focused on the change in the volatilome with regard to each method of anchovies preparation.

Volatile compounds in marinated samples

Integration failed to retrieve all the volatile compounds for each sample. Compounds or sample duplicates with missing values were removed from the dataset. Thus, on the 36 compounds, only 13 were analyzed as shown in **Table 1**. Among the samples, only one sample was

completely removed (*i.e.* both duplicates): a sample preserved for 9 months without any change in storage temperature. For the remaining samples, only four samples had data from only one duplicate.

Figure 1 illustrates the results of the PCA according to storage time. The global analysis (**Figure 1 a**) showed that samples were relatively homogeneous, especially in the first months of storage (from M0 to M4). Groups of sample storage were differentiated primarily along the x-axis. The composition of volatile compounds appeared to change during the first five months of preservation and the product appeared to stabilize in the following months. **Table 2** shows that the samples of the fifth month stand out from the other months with higher proportions of (E,E)-2,4-heptadienal, (E,Z)-2,6-nonadienal, 2,3-octanedione, (E)-2-heptenal, hexanal, heptanal, benzeneacetaldehyde and benzaldehyde. The fifth month may be designated as the month when the volatile compounds reach their optimum with regard to the storage of marinated anchovies.

Based on this five-month optimum, the dataset was then split into a pre-optimal phase and a post-optimal phase. The pre-optimal phase (**Figure 1 b**) shows that months 0 to 4 are characterized by smaller proportions in the above-mentioned compounds compared to the fifth month, except for hexanal and heptanal. Moreover, months 4 and 5 have higher proportions of 2-ethylfuran, (E)-2-hexenal and (E)-2-pentenal than the other months. In the post-optimal phase (**Figure 1 c**), groups were differentiated only along the x-axis. Thus from month 6 to month 9, proportions of 2,3-octanedione, (E,E)-2,4-heptadienal, 2-ethylfuran, (E)-2-hexenal, (E)-2-pentenal, (E)-2-heptenal, hexanal, heptanal and benzeneacetaldehyde decreased.

Splitting the dataset into two phases according to this putative month 5 optimum showed an overall decrease in the proportions of 2-ethylfuran, (E)-2-hexenal and (E)-2-pentenal throughout the study. However, for some compounds, proportions increased up until month 5 before decreasing, e.g. 2,3-octanedione, 2,4-heptadienal, (E)-2-heptenal and benzeneacetaldehyde.

218 Volatile compounds in salted samples

219 For these samples also, integration failed to retrieve all the volatile compounds for each
220 duplicate. Samples with missing values were discarded. Ultimately, data from 24 compounds
221 out to 36 were analyzed as shown in **Table 1**. Five duplicates were discarded due to missing
222 data.

223 **Figure 2** shows the results according to storage time. The storage month groups were
224 differentiated mainly along the x-axis (**Figure 2 a**). Groups were relatively homogeneous
225 especially during the first months of storage (from M0 to M4). This can be observed, in
226 particular, for samples analyzed at the initial stage. A change in the profile of volatile
227 compounds was recorded at six months. Month 6 thus appeared to correspond to the volatile
228 compound optimum for salted anchovies. **Table 2** shows that samples from month 2 stand out
229 with higher proportions of 2-pentenal. Samples from month 6 differ from the other groups,
230 probably due to higher relative quantities in the following compounds: (E,E)-2,4-heptadienal,
231 (Z)-4-heptenal, (E)-2-heptenal, benzeneacetaldehyde, 2-nonanone, (E,E)-3,5-octadien-2-one,
232 heptanal, hexanal, 2,3-octanedione and benzaldehyde. Finally, samples corresponding to the last
233 few months (10, 16 and 18) do not appear different from each other.

234 According to this putative optimum, the dataset was divided into two phases, a pre-optimal
235 phase and a post-optimal phase. The PCA of the pre-optimal phase (**Figure 2 b**) shows that
236 months 0, 2 and 6 were characterized by higher proportions of (E)-2-hexenal, 2-methylbutanal
237 and 3-methylbutanal compared to month 4. In the same phase, the month 6 differs from the
238 other months, showing higher levels for the compounds described for the overall dataset and
239 lower proportions of 1-penten-3-ol. The PCA of the post-optimal phase (**Figure 2 c**) shows that
240 month 8 stands out due to higher levels of 2,3-octanedione, (E,E)-2,4-heptadienal, (E)-2-
241 pentenal, (E,E)-3,5-octadien-2-one, (Z)-4-heptenal, (E)-2-heptenal, benzeneacetaldehyde,

heptanal and hexanal. Moreover, months 10 to 16 show higher relative quantities in TMA. Lastly, there seems to be no differences between month 16 and month 18: they both show higher proportions of 2-methylbutanal, 3-methylbutanal and 2-nonanone compared to previous months.

During the two phases, relative quantities of 2,3-octanedione, (E,E)-2,4-heptadienal, (Z)-4-heptenal, (E)-2-heptenal, (E,E)-3,5-octadien-2-one, hexanal, heptanal and benzeneacetaldehyde appeared to increase until reaching the optimum level before decreasing. During the same period, a continuous increase of 2-nonanone was recorded.

Changes in biogenic amines

The overall results are summarized in **Table 3**. In the large majority of cases, no samples ever exceeded the limit defined by the regulation (Commission Regulation (EC) No 1441/2007 & Commission Regulation (EC) No 1019/2013) during the study. Only marinated anchovies that underwent a break in the cold chain showed a significant change ($p\text{-value} < 0.001$) in the quantity of histamine, exceeding of the prescribed limit from month 6. Regarding putrescine, cadaverine and tyramine, a significant change ($p\text{-value} \leq 0.003$) was also demonstrated with a maximum concentration of nearly 50 mg/kg, between month 4 and 6. Finally a significant change ($p\text{-value} < 0.001$) in the concentrations of spermidine and spermine was recorded, although levels remained low, near 20 mg/kg. The experiment carried out on samples that did not undergo a break in the cold chain shows similar patterns, although all the amine concentrations were lower. Thus the concentrations of histamine, putrescine, cadaverine, tyramine, spermidine and spermine did not exceed the threshold values of 40 mg/kg.

Regarding the salted anchovies, although significant changes were recorded throughout the storage of samples, the threshold of 20 mg/kg was never reached for any amine at any storage time or either storage condition.

DISCUSSION

Principal component analysis helped to visualize a significant proportion of the variance contained in the datasets analyzed. For each analysis, principal components explained at least 64% of the variability in the whole dataset, which is an acceptable level (**Figure 1 & Figure 2**).

In these analyses, a difference in the volatile profile between marinated and salted samples was observed. Generally, it was found that the marinated samples did not show as many compounds as those identified for salted anchovy samples. As a result, multivariate statistical analysis could not be carried out on the same number of compounds for both types of processing methods. A significant amount of acetic acid in the headspace of marinated samples may explain the lower number of compounds recovered, possibly due to saturation of the SPME fiber.

Regarding the nature of compounds allowing the discrimination among samples, a large majority originated from the degradation of fatty acids of the ω -3 and 6 series (**Table 4**). This origin is consistent with observations of increased levels of free fatty acids throughout the maturation process of anchovies.^{3,28} The presence of products of the catabolism of amino acids was also recorded as well as some products such as TMA or 2-ethylfuran, which are frequently observed in fish matrices. All compounds selected for the analysis of marinated samples were used in the statistical analysis of salted anchovy dataset. Among the compounds included in the analysis of salted samples, five come from the degradation of fatty acids, three from the catabolism of amino acids, two from sugar degradation and one from the degradation of trimethylamine N-oxide (TMAO) (**Table 4**).

The PCA traced the changes in the composition of the headspace over time. From those analyses, volatilome optima appear to occur at six months for salted anchovies and five months for marinated anchovies. Only a small change in the composition of volatile compounds was recorded after this period of storage.

290 The results of salted samples corroborate those of a study carried out on the selection of sensory
291 attributes for the evaluation of the maturation of processed anchovies: the smell of Iberian ham
292 is a reliable marker of maturation of salted anchovies.²² SPME-GC/MS studies have specifically
293 shown the presence of heptanal, with a maximum at the sixth month of storage. This compound
294 is described in the literature as having a characteristic odor of ham, fat, even rancidity.^{29,30}
295 Moreover, higher proportions of 2,3-octanedione, 2-nonanone, (Z)-4-heptenal, (E)-2-heptenal,
296 benzaldehyde, hexanal, benzeneacetaldehyde, (E,E)-2,4-heptadienal and (E,E)-3,5-octadien-2-
297 one were recorded for these samples. After six months, the proportions of all compounds
298 decreased except for benzaldehyde. Among the above-mentioned compounds, only 2,3-
299 octanedione and 2-nonanone have never been described in the total volatile profile, in previous
300 studies on ripened anchovy maturation.¹⁹⁻²¹ Based on the aromatic qualifiers defined by Leduc
301 *et al.*¹⁶, odors can be divided into the following classes: earthy/woody smell, vegetable odor and
302 floral smell.³⁰⁻³² Moreover, anchovy samples from the tenth to eighteenth month are
303 characterized by higher proportions of TMA compared to the eighth month. This suggests that
304 the increase in total volatile basic nitrogen (TVB-N), the chemical class to which TMA
305 belongs³³ and recorded during the anchovy maturation process,³ continues during storage.

306 Regarding marinated anchovies, the fifth month appears to mark the volatilome optimum of the
307 product. These samples were characterized by a high proportion of benzeneacetaldehyde, (E,E)-
308 2,4-heptadienal, (E)-2-heptenal, benzaldehyde, (E,Z)-2,6-nonadienal and 2,3-octanedione
309 compared with other months. Only 2,3-octanedione has never been described in other studies on
310 salted anchovies¹⁹⁻²¹ (there are no published studies on the volatilome of marinated anchovies).
311 The proportion of the above-mentioned compounds decreased except benzaldehyde. These
312 compounds are characterized by respective odors of moss/solvent,³² fat/soap,^{34,35}
313 cardboard/fat,^{30,31} almond/woody^{29,30} and cucumber for the latter two.^{31,34}

Five compounds were present in maximum amounts at the volatilome optimum of processed anchovies, whatever the method of preparation: (E,E)-2,4-heptadienal, benzeneacetaldehyde, (E)-2-heptenal, benzaldehyde and 2,3-octanedione. To our knowledge, this is the first time that the latter compound has been identified, only by MS and LRI, as component of volatiles compounds in anchovies. 2-nonanone, hexanal and (E,E)-3,5-octadien-2-one appeared to be more specific to salt-ripened anchovies.

Analysis of biogenic amines showed high variability in quantities in similar samples principally in marinated anchovies stored with rupture in the temperature conditions. This probably reflects the heterogeneity of the fish that make up the samples. The initial microbiological load of the product, hygiene measures and the presence of bruises are all factors that may influence the degradation of anchovies. Despite the high variability of two samples with rupture in the storage conditions, the lower quantities observed exceeded 50 mg.kg^{-1} , concentration where the first allergic symptoms could be observed.

The type of anchovy processing has an impact on the quantity of biogenic amines, because marinated samples had higher concentrations of histamine, cadaverine, putrescine and tyramine than salted anchovies. This fact can be explained by the bacteriostatic action of salt ripening, while the development of lactic flora, potential producers of biogenic amines, is still possible in acidic conditions.²

The major factor that determines the production of biogenic amines, as highlighted in our study and previous papers,^{6,13,14} is storage temperature. If the cold chain is broken during storage of marinated anchovies, the regulatory threshold of 400 mg.kg^{-1} is exceeded. However, it should be noted that the producing process, mainly in salt ripening treatment, limits the bacterial development at the origin of histamine production.⁵²

The present work helps to visualize the changes in the profile of volatile compounds and safety for the both types of anchovy processing techniques. It was shown that irregularities in

339 temperatures of storage lead to the development of biogenic amines in marinated samples,
340 whereas no risk was observed in salted samples. In contrast, the modification of temperature
341 conditions did not impact the volatile profile of marinated or salted samples for which the
342 volatilome optimum is respectively reached after five and six months. Considering these results,
343 none of them lead to conclude to a food safety risk caused by biogenic amines when anchovies
344 are stored in good temperature conditions. However, the study of volatilomes suggests that
345 marinated and salted anchovies should be consumed within five to six months after maturation,
346 respectively, so as to guarantee a product with optimal flavor for consumers.

ABBREVIATIONS USED

ALA: α -Linolenic acid, CAR/PDMS: Carboxen/polydimethylsiloxane, DHA: Docosahexaenoic acid, EPA: Eicosapentaenoic acid, GC: Gas chromatography, HACCP: Hazard analysis critical control point, HPLC: High performance liquid chromatography, LA: Linoleic acid, LRI: Linear retention index, MS: Mass spectrometry, MUFA: Mono-unsaturated fatty acid, PCA: Principal component analysis, PUFA : Poly-unsaturated fatty acid, SPME: Solid phase microextraction, TMA: Trimethylamine, TMAO: Trimethylamine N-oxide, TVB-N: Total volatile basic nitrogen, wR: with rupture & w/oR: without rupture.

ACKNOWLEDGEMENTS

We thank Assi N'Guessan, Ossarath Kol and Philippe Droin for their critical reviews of this article and Ophélie Frottier and Paul Morieux for their kind technical assistance. Funds for this study, for which the CITPPM was project leader, were provided by FranceAgriMer, run by the French Ministry of Agriculture, as part of the programme "*Soutien à l'innovation dans la filière des produits de la pêche et de l'aquaculture – Edition 2010*". A. Dehaut is grateful to the Nord-Pas de Calais Regional Council (France) and ANSES for providing financial support for his PhD studies.

REFERENCES

1. FAO, The state of world fisheries and aquaculture 2012. In Food and Agriculture Organization, Ed. FAO: 2012; p 24.
2. Fuselli, S.; Casales, M.; Fritz, R.; Yeannes, M., Microbiology of the marination process used in anchovy (*Engraulis anchoita*) production. *LWT-Food Sci. Tech.* **1994**, *27*, 214-218.
3. Hernández-Herrero, M. M.; Roig-Sagués, A. X.; López, S. E. I.; Rodríguez-Jerez, J. J.; Mora-Ventura, M. T., Total volatile basic nitrogen and other physico-chemical and microbiological characteristics as related to ripening of salted anchovies. *J. Food Sci.* **1999**, *64*, 344-347.
4. El-Filali, F.; Hanoune, S.; Khbaya, B.; M'Handi, N.; Kaaya, A. In *Histamine and microbiological change during the storage of semi-preserved anchovies*, Second workshop on fish technology, utilization and quality assurance in Africa, Agadir, Morocco, 24-28 November 2008., 2009; Food and Agriculture Organization of the United Nations (FAO): 2009; pp 133-137.
5. Rodríguez-Jerez, J.; Lopez-Sabater, E.; Hernandez-Herrero, M.; Mora-Ventura, M., Histamine, putrescine and cadaverine formation in Spanish semipreserved anchovies as affected by time/temperature. *J. Food Sci.* **1994**, *59*, 993-997.
6. Karaçam, H.; Kutlu, S.; Köse, S., Effect of salt concentrations and temperature on the quality and shelf-life of brined anchovies. *Int. J. Food Sci. Technol.* **2002**, *37*, 19-28.
7. Sen, M.; Temelli, S., Microbiological and chemical qualities of marinated anchovy prepared with different vegetable additives and sauce. *Rev. Méd. Vét.* **2003**, *154*, 703-708.
8. EFSA Panel on Biological Hazards (BIOHAZ), Scientific opinion on risk based control of biogenic amine formation in fermented foods. *EFSA Journal* **2011**, *9*, 2393.
9. Commission Regulation (EC) N° 1441/2007 of 5 December 2007 amending Regulation (EC) N° 2073/2005 on microbiological criteria for foodstuffs. *Off. J. Eur. Union* **2007**, *322*, 1-18.
10. Commission Regulation (EC) N° 1019/2013 of 23 October 2013 amending Annex I to Regulation (EC) No 2073/2005 as regards histamine in fishery products. *Off. J. Eur. Union* **2013**, *282*, 46-47.
11. Pons-Sanchez-Cascado, S.; Veciana-Nogues, M.; Bover-Cid, S.; Marine-Font, A.; Vidal-Carou, M., Volatile and biogenic amines, microbiological counts, and bacterial amino acid decarboxylase activity throughout the salt-ripening process of anchovies (*Engraulis encrasicolus*). *J. Food Prot.* **2005**, *68*, 1683-1689.
12. Pons-Sanchez-Cascado, S.; Vidal-Carou, M. C.; Marine-Font, A.; Veciana-Nogues, M. T., Influence of the freshness grade of raw fish on the formation of volatile and biogenic amines during the manufacture and storage of vinegar-marinated anchovies. *J. Agric. Food Chem.* **2005**, *53*, 8586-8592.
13. Veciana-Nogues, M.; Marine-Font, A.; Vidal-Carou, M., Changes in biogenic amines during the storage of Mediterranean anchovies immersed in oil. *J. Agric. Food Chem.* **1997**, *45*, 1385-1389.
14. Veciana-Nogues, M.; Vidal-Carou, M.; Marine-Font, A., Histamine and tyramine in preserved and semi-preserved fish products. *J. Food Sci.* **1989**, *54*, 1653-1655.
15. Béné, A.; Hayman, A.; Reynard, E.; Luisier, J. L.; Villettaz, J. C., A new method for the rapid determination of volatile substances: the SPME-direct method: Part II. Determination of the freshness of fish. *Sensor Actuat B-Chem.* **2001**, *72*, 204-207.
16. Leduc, F.; Tournayre, P.; Kondjoyan, N.; Mercier, F.; Malle, P.; Kol, O.; Berdagué, J. L.; Duflos, G., Evolution of volatile odorous compounds during the storage of European seabass (*Dicentrarchus labrax*). *Food Chem.* **2012**, *131*, 1304-1311.

17. Duflos, G.; Leduc, F.; N'Guessan, A.; Krzewinski, F.; Kol, O.; Malle, P., Freshness characterisation of whiting (*Merlangius merlangus*) using an SPME/GC/MS method and a statistical multivariate approach. *J. Sci. Food Agric.* **2010**, *90*, 2568-2575.
18. Leduc, F.; Krzewinski, F.; Le Fur, B.; N'Guessan, A.; Malle, P.; Kol, O.; Duflos, G., Differentiation of fresh and frozen/thawed fish, European sea bass (*Dicentrarchus labrax*), gilthead seabream (*Sparus aurata*), cod (*Gadus morhua*) and salmon (*Salmo salar*), using volatile compounds by SPME/GC/MS. *J. Sci. Food Agric.* **2012**, *92*, 2560-2568.
19. Triqui, R.; Reineccius, G. A., Changes in flavor profiles with ripening of anchovy (*Engraulis encrasicolus*). *J. Agric. Food Chem.* **1995**, *43*, 1883-1889.
20. Triqui, R.; Reineccius, G. A., Flavor development in the ripening of anchovy (*Engraulis encrasicolus* L.). *J. Agric. Food Chem.* **1995**, *43*, 453-458.
21. Triqui, R.; Guth, H., Determination of potent odorants in ripened anchovy (*Engraulis encrasicolus* L) by aroma extract dilution analysis and by gas chromatography-olfactometry of headspace samples. In *Flavor and Lipid Chemistry of Seafoods*, American Chemical Society: 1997; Vol. 674, pp 31-38.
22. Besteiro, I.; Rodriguez, C. J.; Tilve-Jar, C.; Pascual, C., Selection of attributes for the sensory evaluation of anchovies during the ripening process. *J. Sens. Stud.* **2000**, *15*, 65-77.
23. Kondjoyan, N.; Berdagué, J.-L., *A compilation of relative retention indices for the analysis of aromatic compounds*. Ed. du Laboratoire Flaveur: 1996.
24. Van den Dool, H.; Dec Kratz, P., A generalization of the retention index system including linear temperature programmed gas—liquid partition chromatography. *J. Chromatogr., A* **1963**, *11*, 463-471.
25. Malle, P.; Valle, M.; Bouquelet, S., Assay of biogenic amines involved in fish decomposition. *J. AOAC Int.* **1995**, *79*, 43-49.
26. Duflos, G.; Dervin, C.; Malle, P.; Bouquelet, S., Relevance of matrix effect in determination of biogenic amines in plaice (*Pleuronectes platessa*) and whiting (*Merlangus merlangus*). *J. AOAC Int.* **1999**, *82*, 1097-1101.
27. Rakotomalala, R., TANAGRA : un logiciel gratuit pour l'enseignement et la recherche. In *Extraction et Gestion des Connaissances EGC 2005*, Cépaduès-Editions, Ed. Toulouse, 2005; Vol. 2, pp 697-702.
28. Roldan, H. A.; Barassi, C. A.; Trucco, R. E., Increase on free fatty acids during ripening of anchovies (*Engraulis anchoita*). *Int. J. Food Sci. Technol.* **1985**, *20*, 581-585.
29. García-González, D. L.; Tena, N.; Aparicio-Ruiz, R.; Morales, M. T., Relationship between sensory attributes and volatile compounds qualifying dry-cured hams. *Meat Sci.* **2008**, *80*, 315-325.
30. Giri, A.; Osako, K.; Okamoto, A.; Ohshima, T., Olfactometric characterization of aroma active compounds in fermented fish paste in comparison with fish sauce, fermented soy paste and sauce products. *Food Res. Int.* **2010**, *43*, 1027-1040.
31. Ólafsdóttir, G.; Fleurence, J., Evaluation of fish freshness using volatile compounds - Classification of volatile compounds in fish. In *Methods to determine the freshness of fish in research and industries*, Proceedings of the Final Meeting of the Concerted Action (Evaluation of Fish Freshness), Nantes, France, Nov 12-14 1997, Institut International du Froid Publication, Paris, 1997.
32. Varlet, V.; Knockaert, C.; Prost, C.; Serot, T., Comparison of odor-active volatile compounds of fresh and smoked salmon. *J. Agric. Food Chem.* **2006**, *54*, 3391-3401.
33. Gill, T. A., Objective analysis of seafood quality. *Food Reviews International* **1990**, *6*, 681-714.

34. Triqui, R.; Bouchriti, N., Freshness assessments of moroccan sardine (*Sardina pilchardus*) : comparison of overall sensory changes to instrumentally determined volatiles. *J. Agric. Food Chem.* **2003**, *51*, 7540-7546.
35. Jónsdóttir, R. s.; Sveinsdóttir, K. n.; Magnússon, H.; Arason, S. n.; Lauritzen, K.; Thorarinsdóttir, K. A., Flavor and quality characteristics of salted and desalted cod (*Gadus morhua*) produced by different salting methods. *J. Agric. Food Chem.* **2011**, *59*, 3893-3904.
36. Duflos, G.; Coin, V. M.; Cornu, M.; Antinelli, J.-F.; Malle, P., Determination of volatile compounds to characterize fish spoilage using headspace/mass spectrometry and solid-phase microextraction/gas chromatography/mass spectrometry. *J. Sci. Food Agric.* **2006**, *86*, 600-611.
37. Sinz, Q.; Schwab, W., Metabolism of amino acids, dipeptides and tetrapeptides by *Lactobacillus sakei*. *Food Microbiol.* **2012**, *29*, 215-223.
38. Yaylayan, V. A.; Keyhani, A., Origin of 2,3-pentanedione and 2,3-butanedione in d-glucose/l-alanine maillard model Systems. *J. Agric. Food Chem.* **1999**, *47*, 3280-3284.
39. Álvarez-Martín, P.; Belén Flórez, A.; Hernández-Barranco, A.; Mayo, B., Interaction between dairy yeasts and lactic acid bacteria strains during milk fermentation. *Food Control* **2008**, *19*, 62-70.
40. McSweeney, P. L. H.; Sousa, M. J., Biochemical pathways for the production of flavour compounds in cheeses during ripening: A review. *Lait* **2000**, *80*, 293-324.
41. Jaffrès, E.; Lalanne, V.; Macé, S.; Cornet, J.; Cardinal, M.; Sérot, T.; Dousset, X.; Joffraud, J.-J., Sensory characteristics of spoilage and volatile compounds associated with bacteria isolated from cooked and peeled tropical shrimps using SPME–GC–MS analysis. *Int. J. Food Microbiol.* **2011**, *147*, 195-202.
42. Kawai, T.; Sakaguchi, M., Fish flavor. *Crit. Rev. Food Sci. Nutr.* **1996**, *36*, 257-298.
43. Hsieh, R. J.; Kinsella, J. E., *Oxidation of polyunsaturated fatty acids : Mechanisms, Products, and Inhibition with emphasis on Fish*. Academic Press, Inc.: 1989; Vol. 33, p 109.
44. Josephson, D. B.; Lindsay, R. C., Retro-aldol related degradations of 2,4-decadienal in the development of staling flavors in fried foods. *J. Food Sci.* **1987**, *52*, 1186-1190.
45. Iglesias, J.; Medina, I., Solid-phase microextraction method for the determination of volatile compounds associated to oxidation of fish muscle. *J. Chromatogr., A* **2008**, *1192*, 9-16.
46. Medina, I.; Satué-Gracia, M.; Frankel, E., Static headspace gas chromatographic analyses to determine oxidation of fish muscle lipids during thermal processing. *J. Am. Oil Chem. Soc.* **1999**, *76*, 231-236.
47. Grosch, W., Reactions of hydroperoxides - products of low molecular weight. In *Autoxidation of Unsaturated Lipids*, Chan, H. W. S., Ed. Academic Press London, 1987; pp 95-139.
48. Meynier, A.; Genot, C.; Gandemer, G., Volatile compounds of oxidized pork phospholipids. *J. Am. Oil Chem. Soc.* **1998**, *75*, 1-7.
49. Young, O. A.; Berdagué, J. L.; Viallon, C.; Rousset-Akrim, S.; Theriez, M., Fat-borne volatiles and sheepmeat odour. *Meat Sci.* **1997**, *45*, 183-200.
50. Frankel, E. N., Volatile lipid oxidation products. *Prog. Lipid Res.* **1983**, *22*, 1-33.
51. Frankel, E., *Lipid Oxidation*. Second Edition ed.; The Oily Press Lipid Library: 2005; Vol. 18, p 470.
52. Nei, D., Evaluation of Non-bacterial factors contributing to histamine accumulation in fish fillets. *Food Control* **2014**, *35*, 142-145.

FIGURE CAPTIONS

Figure 1: PCA obtained on the marinated anchovy samples for the whole dataset (a), from the pre-optimal phase (b) and from the post-optimal phase (c). Symbols indicate the tested month (M) of storage: (M0 ●, M2 ●, M3 ●, M4 ●, M5 ●, M6 ■, M7 ■, M8 ■ and M9 ■).

Figure 2: PCA obtained on the salted samples for the whole dataset (a), from the pre-optimal phase (b) and from the post-optimal phase (c). Symbols indicate the tested month (M) of storage (M0 ●, M2 ●, M4 ●, M6 ●, M8 ■, M10 ■, M16 ■ and M18 ■).

TABLES

Table 1: Settings used to perform automatic integration of the chromatograms.

compound ^a	LRI ^d	reliability ^e	ions used for integration (m/z) ^f
ethanol ^c	477	MS, LRI, Std	45 (46 43)
dimethyl sulfide	516	MS, LRI, Std	62 (47 45 46 61)
trimethylamine ^c	522	tentative	58 (59 42)
2-methyl- propanal ^c	549	MS, LRI, Std	43 (41 72)
2,3-butanedione ^c	586	MS, LRI, Std	43 (86)
butanal	591	MS, LRI, Std	44 (43 72 57 41)
2-butanone ^{b,c}	595	Tentative	43 (72 57)
3-methyl- butanal ^c	652	MS, LRI, Std	41 (44 58 43 39)
2-methyl- butanal ^c	662	MS, LRI, Std	41 (57 58 39)
1-penten-3-ol ^c	684	MS, LRI, Std	57 (41 43 39 58)
2,3-pentanedione	697	Tentative	43 (57 100)
2-ethyl-furan ^{b,c}	700	MS, LRI, Std	81 (96 53)
3-methyl-1-butanol	753	Tentative	55 (42 43 70)
(E)-2-pentenal ^{b,c}	755	Tentative	55 (84 83 41)
1-pentanol	768	MS, LRI, Std	42 (55 41 70)
2,3-hexanedione	785	MS, LRI, Std	43 (41 71 57)
hexanal ^{b,c}	799	MS, LRI, Std	44 (56 41 43 72 82)
(E)-2-hexenal ^{b,c}	855	Tentative	41 (42 39 83 69)
(Z)-4-heptenal ^c	898	Tentative	41 (68 55 84 43)
heptanal ^{b,c}	900	MS, LRI, Std	70 (41 55 57 81 44)
6-methyl-2-heptanone	956	Tentative	43 (58 71 95 110 41)
(E)-2-heptenal ^{b,c}	960	MS, LRI, Std	41 (68 55 84 43)
benzaldehyde ^{b,c}	970	MS, LRI, Std	77 (106 105 51)
1-octen-3-ol	981	MS, LRI, Std	57 (43 72 55 85)
2,3-octanedione ^{b,c}	983	Tentative	43 (71 99)
(E,E)-2,4-heptadienal [1] ^{b,c}	997	MS, LRI, Std	81 (110 53 41 67 79)
octanal	1002	MS, LRI, Std	43 (44 56 84 69 100)
(E,E)-2,4-heptadienal [2] ^{b,c}	1014	tentative	81 (110 53 41 67 79)
benzeneacetaldehyde ^{b,c}	1053	MS, LRI, Std	91 (120 92 65)
(E)-2-octenal	1064	Tentative	70 (55 41 83 57 69)
(E,E)-3,5-octadien-2-one [1] ^c	1073	tentative	95 (43 81 109 124)
2-nonanone ^c	1091	Tentative	43 (58 71 57 41)
(E,E)-3,5-octadien-2-one [2] ^c	1096	Tentative	95 (43 81 109 124)
nonanal	1106	MS, LRI, Std	57 (41 56 70 82 98)
(E,Z)-2,6-nonadienal ^{b,c}	1159	MS, LRI, Std	41 (69 70 53)
decanal	1224	MS, LRI, Std	43 (57 70 82 95 112)

^a Numbers in brackets correspond to identifiers of peaks when a double peak is recorded on chromatograms for one identified compound. ^b Compounds used for multivariate analysis of marinated samples. ^c Compounds used for multivariate analysis of salted samples. ^d Linear retention index (LRI). ^e MS, LRI, Std: identification confirmed by mass spectrometry, LRI and pure standard; Tentative: tentative identification by mass spectrometry and LRI; tentative: tentatively identified compounds by mass spectrometry only. ^f **43**: target ion for the method. (41 72): ions of reference to improve the integration process.

Table 2: Correlation coefficients between chemical compounds and the two first axes determined by PCA.

Three PCA were carried out per maturation process.

Marinated anchovies			
	Whole dataset	Pre-optimum phase	Post-optimum phase
PC 1	benzaldehyde (0.93) ^a	benzaldehyde (0.95)	(E,E)-2,4-heptadienal [2] (0.96)
	2,3-octanedione (0.88)	(E)-2-heptenal (0.93)	heptanal (0.95)
	(E,E)-2,4-heptadienal [2] ^b (0.86)	2,3-octanedione (0.9)	hexanal (0.83)
	(E)-2-heptenal (0.85)	(E,Z)-2,6-nonadienal (0.86)	(E)-2-heptenal (0.83)
	benzeneacetaldehyde (0.83)	(E,E)-2,4-heptadienal [2] (0.81)	(E)-2-hexenal (0.83)
	heptanal (0.82)	benzeneacetaldehyde (0.78)	(E)-2-pentenal (0.8)
	(E,Z)-2,6-nonadienal (0.75)	(E,E)-2,4-heptadienal [1] (0.75)	2,3-octanedione (0.76)
	hexanal (0.71)		2-ethyl-furan (0.74)
			benzaldehyde (0.71)
			(E,E)-2,4-heptadienal [1] (0.74)
(Inertia)	(49.3%) ^c	(48.8%)	(56.4%)
PC 2	(E)-2-pentenal (0.92)	(E)-2-hexenal (-0.95)	(E,Z)-2,6-nonadienal (-0.9)
	2-ethyl-furan (0.92)	(E)-2-pentenal (-0.91)	
	(E)-2-hexenal (0.84)	2-ethyl-furan (-0.9)	
(Inertia)	(27.9%)	(29.9%)	(25.6%)
Salted anchovies			
	Whole dataset	Pre-optimum phase	Post-optimum phase
PC 1	(E,E)-3,5-octadien-2-one [1] (0.97)	(E,E)-3,5-octadien-2-one [1] (0.98)	(E,E)-3,5-octadien-2-one [2] (0.95)
	(Z)-4-heptenal (0.96)	heptanal (0.97)	(E,E)-3,5-octadien-2-one [1] (0.93)
	(E,E)-3,5-octadien-2-one [2] (0.95)	2-nonanone (0.97)	(E,E)-2,4-heptadienal [1] (0.91)
	heptanal (0.95)	(Z)-4-heptenal (0.97)	(E,E)-2,4-heptadienal [2] (0.91)
	(E)-2-heptenal (0.94)	(E,E)-3,5-octadien-2-one [2] (0.97)	(Z)-4-heptenal (0.89)
	hexanal (0.94)	(E)-2-heptenal (0.95)	(E)-2-heptenal (0.85)
	2-nonanone (0.92)	hexanal (0.95)	(E)-2-pentenal (0.82)
	(E,E)-2,4-heptadienal [2] (0.92)	benzaldehyde (0.92)	benzeneacetaldehyde (0.78)
	2,3-octanedione (0.87)	(E,E)-2,4-heptadienal [2] (0.9)	hexanal (0.75)
	benzaldehyde (0.85)	benzeneacetaldehyde (0.9)	heptanal (0.73)
	(E,E)-2,4-heptadienal [1] (0.79)	2,3-octanedione (0.87)	2,3-octanedione (0.71)
	benzeneacetaldehyde (0.77)	(E,E)-2,4-heptadienal [1] (0.72)	trimethylamine (-0.84)
		1-penten-3-ol (-0.7)	
(Inertia)	(52.9%)	(57.7%)	(45.4%)
PC 2	(E)-2-pentenal (-0.83)	3-methyl-butanal (-0.78)	3-methyl-butanal (0.85)
		2-methyl-butanal (-0.74)	2-nonanone (0.76)
		(E)-2-hexenal (-0.72)	2-methyl-butanal (0.73)
(Inertia)	(12.0%)	(15.2%)	(18.2%)

^a Coefficient in parenthesis correspond to the correlation of the volatile compound with the PC concerned (row) for a given PCA (column). ^b Numbers in brackets correspond to identifiers of peaks when a double peak is recorded on chromatograms for one identified compound. ^c Percentage in parenthesis correspond to the inertia harbored by the PC concerned for a given PCA (e.g 49.3% means that PC1 harbored 49.3% of the dataset inertia in the PCA achieved on the marinated anchovies whole dataset).

Table 3: Changes in the concentrations of biogenic amines throughout a shelf-life of 14 months^a. Values are listed according to curing method and storage conditions with (wR) or without (w/oR) rupture in the temperature of storage.

	histamine (mg.kg ⁻¹) ^b				putrescine (mg.kg ⁻¹)			
	marinated		salted		marinated		salted	
	wR	w/oR	wR	w/oR	wR	w/oR	wR	w/oR
M0	4 ± 1 a	4 ± 1 a	1 ± 1 a	1 ± 1	4 ± 1 a	4 ± 1 a	3 ± 1 a	3 ± 1 a
M2	11 ± 2 b	6 ± 1 a	n.d. ^d	2 ± 2	4 ± 0 a	3 ± 1 a	2 ± 2 a	3 ± 2 ab
M4	28 ± 15 b	21 ± 11 b	1 ± 2 a	1 ± 2	25 ± 14 b	17 ± 8 b	3 ± 1 a	2 ± 0 ab
M6	2826 ± 2356 c	4 ± 2 a	n.d.	n.d.	48 ± 13 b	4 ± 0 a	7 ± 1 b	7 ± 1 b
M8	369 ± 269 c	7 ± 4 ab	n.d.	n.d.	21 ± 6 b	4 ± 1 a	3 ± 1 a	4 ± 0 b
M10			n.d.	n.d.			6 ± 4 ab	5 ± 1 ab
M12			1 ± 1 a	n.d.			8 ± 1 b	6 ± 1 b
M14			1 ± 1 a	n.d.			4 ± 0 a	4 ± 1 ab
<i>p</i> ^c	< 0.001	0.003	0.007	0.291	< 0.001	0.002	< 0.001	< 0.001
	tyramine (mg.kg ⁻¹)				cadaverine (mg.kg ⁻¹)			
	marinated		salted		marinated		salted	
	wR	w/oR	wR	w/oR	wR	w/oR	wR	w/oR
M0	14 ± 3 a	14 ± 3 a	7 ± 2 a	7 ± 2 a	15 ± 2 a	15 ± 2 a	12 ± 2 a	12 ± 2 a
M2	4 ± 0 b	5 ± 2 b	7 ± 1 a	7 ± 1 a	n.d.	12 ± 1 a	2 ± 1 b	2 ± 2 ab
M4	38 ± 25 a	20 ± 11 b	4 ± 1 a	3 ± 1 a	48 ± 23 b	36 ± 13 b	3 ± 0 b	4 ± 1 ab
M6	6 ± 3 b	2 ± 1 a	10 ± 1 a	9 ± 1 a	36 ± 8 b	38 ± 1 b	7 ± 1 ab	8 ± 1 ab
M8	10 ± 9 ab	3 ± 2 b	5 ± 1 a	4 ± 0 a	14 ± 4 a	14 ± 1 a	4 ± 1 b	3 ± 0 b
M10			8 ± 2 a	5 ± 1 a			7 ± 2 ab	5 ± 1 ab
M12			10 ± 2 a	5 ± 0 a			6 ± 1 ab	4 ± 1 b
M14			5 ± 0 a	4 ± 0 a			4 ± 0 ab	3 ± 0 b
<i>p</i>	0.003	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
	spermidine (mg.kg ⁻¹)				spermine (mg.kg ⁻¹)			
	marinated		salted		marinated		salted	
	wR	w/oR	wR	w/oR	wR	w/oR	wR	w/oR
M0	10 ± 3 a	10 ± 3 a	18 ± 3 a	18 ± 3 a	2 ± 1 a	2 ± 1 a	4 ± 1 a	4 ± 1 a
M2	22 ± 2 b	20 ± 5 b	15 ± 4 ab	14 ± 3 ab	13 ± 2 b	12 ± 4 b	11 ± 4 ab	11 ± 3 b
M4	8 ± 3 a	13 ± 4 ab	9 ± 4 ab	13 ± 1 ab	6 ± 3 ab	8 ± 2 b	3 ± 0 ab	2 ± 1 ab
M6	7 ± 3 a	10 ± 1 a	8 ± 1 b	8 ± 1 b	n.d.	1 ± 1 a	n.d.	n.d.
M8	15 ± 3 ab	11 ± 1 a	8 ± 1 b	7 ± 1 b	1 ± 1 a	5 ± 2 b	2 ± 2 b	4 ± 1 ab
M10			11 ± 2 ab	9 ± 1 ab			n.d.	n.d.
M12			9 ± 2 b	8 ± 0 b			n.d.	2 ± 1 ab
M14			8 ± 0 b	7 ± 2 b			3 ± 0 b	4 ± 0 ab
<i>p</i>	< 0.001	0.013	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001

^a Means with different letters are significantly different ($p < 0.05$) throughout the tested period. ^b Means of the concentrations ± 2 standard-error of the mean (95% confidence interval). ^c *p* value calculated with the Kruskal-Wallis test. ^d n.d.: values below the limit of quantitation.

Table 4: Compounds used in the multivariate statistical analysis of anchovies and their assumed origins.

Compounds	Assumed origin
ethanol	Fermentation of sugar
trimethylamine	Trimethylamine-N oxide (TMAO) reduction, degradation of choline ³⁶
2-methyl-propanal	Amino-acid catabolism (L-valine) ³⁷
2,3-butanedione	Fermentation of sugar ^{38,39}
2-butanone	Degradation of 2,3-butanedione ⁴⁰
3-methyl-butanal	Amino-acid catabolism (L-leucine) ³⁷
2-methyl-butanal	Amino-acid catabolism ⁴¹
1-penten-3-ol	Eicosapentaenoic acid (EPA) (ω -3) oxidation ⁴²
(E)-2-pentenal	α -Linolenic acid (ALA) (ω -3) oxidation ⁴³
hexanal	Linoleic acid (LA) (ω -6) oxidation and 2,4-decadienal degradation ⁴³⁻⁴⁵
2-ethyl-furan	ω -3 fatty acid oxidation (ALA, EPA, Docosahexaenoic acid (DHA)) ⁴⁶
(E)-2-hexenal	EPA (ω -3) oxidation ⁴²
(Z)-4-heptenal	EPA (ω -3) oxidation ⁴²
(E)-2-heptenal	Linoleic acid (LA) (ω -6) oxidation ⁴⁷
heptanal	Auto-oxidation of MUFA and PUFA (ω -6) ^{36,48}
benzaldehyde	Amino-acid catabolism ^{37,40}
2,3-octanedione	Lipid oxidation ⁴⁹
(E,E)-2,4-heptadienal	EPA and ALA (ω -3) oxidation ^{42,50}
benzeneacetaldehyde	Amino-acid catabolism ⁴⁰
2-nonanone	Lipid auto-oxidation
(E,E)-3,5-octadien-2-one	EPA (ω -3) auto-oxidation ⁴²
(E,Z)-2,6-nonadienal	ω -3fatty acid oxidation (ALA, EPA, DHA) ⁵¹

FIGURES

Figure 1

Figure 2

TOC GRAPHIC