



**HAL**  
open science

## **Mycobacterium microti infection in a cow in France**

Lorraine Michelet, Krystel de Cruz, Claudine Karoui, Sylvie Hénault, Maria  
Laura Boschioli

► **To cite this version:**

Lorraine Michelet, Krystel de Cruz, Claudine Karoui, Sylvie Hénault, Maria Laura Boschioli. Mycobacterium microti infection in a cow in France. Veterinary Record, 2017, 180, pp.429 - 429. 10.1136/vr.j2041 . anses-03968093

**HAL Id: anses-03968093**

**<https://anses.hal.science/anses-03968093>**

Submitted on 13 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

## ***Mycobacterium microti* infection in a cow in France**

Lorraine Michelet, Krystel de Cruz, Claudine Karoui, Sylvie Hénault, Maria Laura Boschioli

National Reference Laboratory for Tuberculosis, Bacterial Zoonosis Unit, Animal Health Laboratory, ANSES, Maisons-Alfort, France

Corresponding author: [maria-laura.boschioli@anses.fr](mailto:maria-laura.boschioli@anses.fr)

*Mycobacterium microti* is a member of the *Mycobacterium tuberculosis* complex (MTBC), which also includes major human and animal pathogens such as *M. tuberculosis* and *M. bovis*. *M. microti* was originally described as the cause of tuberculosis in wild rodents, but it can also infect several other animal species including people, highlighting its zoonotic potential (Cavanagh and others 2002; Panteix and others 2010).

In 2015, we reported a case of *M. microti* infection in a dairy goat, showing that this mycobacterium could be responsible for misleading bovine tuberculosis (bTB) diagnostic results (Michelet and others 2016). An extended investigation in the same region led to the discovery of an additional case in a cow. As for the goat herd, the suspected cattle herd shared the same pastures with two bTB (*M. bovis*) confirmed herds identified in 2005 and 2015. As a result, the cattle herd was submitted to the single intradermal cervical comparative test. One animal with a doubtful result was culled for direct diagnosis. No lesions were observed at the abattoir on mediastinal, retropharyngeal and tracheobronchial lymph nodes. Samples were analysed and submitted to bacterial culture and PCR (Courcoul and others 2014). Culture was negative after three months in all samples. However a doubtful result was obtained by PCR on the retropharyngeal lymph node. Further characterisation was performed on the extracted DNA employing molecular tools targeting the Regions of Difference which allow differentiation of the MTBC members (Huard and others 2003) and through spoligotyping (Zhang and others 2010). All these tests enabled the identification of *M. microti* spoligotype SB0118, which is the same genotype identified in the goat and in a badger of the same region.

*M. microti* had previously been isolated in skin test reactor cattle in the UK (Jahans and others 2004), demonstrating the risk of infection in livestock and of interference with bTB diagnosis. All these findings raise concern about the reliability of diagnostic tests used for bTB surveillance. The use of highly specific tests based on specific antigens such as ESAT 6 and CFP10 (which are absent in *M. microti* and already currently used in the interferon gamma test employed in France (Faye and others 2011)) are highly recommended to accurately identify *M. bovis* (or *M. caprae*) infection at ante-mortem examination. At post-mortem diagnosis, the use of specific molecular tools should be considered to rapidly distinguishing pathogens within the MTBC and avoid misleading diagnosis.

## References

- CAVANAGH, R., BEGON, M., BENNETT, M., ERGON, T., GRAHAM, I. M., DE HAAS, P. E., HART, C. A., KOEDAM, M., KREMER, K., LAMBIN, X., ROHOLL, P. & SOOLINGEN DV, D. (2002) *Mycobacterium microti* infection (vole tuberculosis) in wild rodent populations. *Journal of clinical microbiology* 40, 3281-3285
- COURCOUL, A., MOYEN, J. L., BRUGERE, L., FAYE, S., HENAULT, S., GARES, H. & BOSCHIROLI, M. L. (2014) Estimation of sensitivity and specificity of bacteriology, histopathology and PCR for the confirmatory diagnosis of bovine tuberculosis using latent class analysis. *PLoS One* 9, e90334
- FAYE, S., MOYEN, J. L., GARES, H., BENET, J. J., GARIN-BASTUJI, B. & BOSCHIROLI, M. L. (2011) Determination of decisional cut-off values for the optimal diagnosis of bovine tuberculosis with a modified IFN $\gamma$  assay (Bovigam(R)) in a low prevalence area in France. *Veterinary microbiology* 151, 60-67
- HUARD, R. C., LAZZARINI, L. C., BUTLER, W. R., VAN SOOLINGEN, D. & HO, J. L. (2003) PCR-based method to differentiate the subspecies of the *Mycobacterium tuberculosis* complex on the basis of genomic deletions. *Journal of clinical microbiology* 41, 1637-1650
- JAHANS, K., PALMER, S., INWALD, J., BROWN, J. & ABAYAKOON, S. (2004) Isolation of *Mycobacterium microti* from a male Charolais-Hereford cross. *The Veterinary record* 155, 373-374
- MICHELET, L., DE CRUZ, K., PHALENTE, Y., KAROUI, C., HENAULT, S., BERAL, M. & BOSCHIROLI, M. L. (2016) *Mycobacterium microti* Infection in Dairy Goats, France. *Emerging infectious diseases* 22, 569-570
- PANTEIX, G., GUTIERREZ, M. C., BOSCHIROLI, M. L., ROUVIERE, M., PLAIDY, A., PRESSAC, D., PORCHERET, H., CHYDERIOTIS, G., PONSADA, M., VAN OORTEGEM, K., SALLOUM, S., CABUZEL, S., BANULS, A. L., VAN DE PERRE, P. & GODREUIL, S. (2010) Pulmonary tuberculosis due to *Mycobacterium microti*: a study of six recent cases in France. *Journal of medical microbiology* 59, 984-989
- ZHANG, J., ABADIA, E., REFREGIER, G., TAJAJ, S., BOSCHIROLI, M. L., GUILLARD, B., ANDREMONT, A., RUIJMY, R. & SOLA, C. (2010) *Mycobacterium tuberculosis* complex CRISPR genotyping: improving efficiency, throughput and discriminative power of 'spoligotyping' with new spacers and a microbead-based hybridization assay. *Journal of medical microbiology* 59, 285-294