



**HAL**  
open science

## Draft genome sequence of a CTX-M-15-producing Escherichia coli ST345 from commercial chicken meat in Brazil

Tiago Casella, Louise T Cerdeira, Miriam R Fernandes, Tiago A Souza,  
Marisa Haenni, Jean-Yves Madec, Nilton Lincopan, Mara C L Nogueira

### ► To cite this version:

Tiago Casella, Louise T Cerdeira, Miriam R Fernandes, Tiago A Souza, Marisa Haenni, et al.. Draft genome sequence of a CTX-M-15-producing Escherichia coli ST345 from commercial chicken meat in Brazil. *Journal of Global Antimicrobial Resistance*, 2017, 9, pp.124 - 125. 10.1016/j.jgar.2017.04.002 . anses-04013271

**HAL Id: anses-04013271**

**<https://hal-anses.archives-ouvertes.fr/anses-04013271>**

Submitted on 3 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright



## Genome Note

Draft genome sequence of a CTX-M-15-producing *Escherichia coli* ST345 from commercial chicken meat in Brazil

Tiago Casella<sup>a</sup>, Louise T. Cerdeira<sup>b</sup>, Miriam R. Fernandes<sup>b</sup>, Tiago A. Souza<sup>c</sup>,  
Marisa Haenni<sup>d</sup>, Jean-Yves Madec<sup>d</sup>, Nilton Lincopan<sup>b,e</sup>, Mara C.L. Nogueira<sup>f,\*</sup>

<sup>a</sup> Instituto de Biociências, Letras e Ciências Exatas, Universidade Estadual Paulista 'Júlia de Mesquita Filho', São José do Rio Preto, Brazil

<sup>b</sup> Departamento de Análises Clínicas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, Brazil

<sup>c</sup> Genome Investigation and Analysis Laboratory (GENIAL), Institute of Biomedical Sciences, Universidade de São Paulo, São Paulo, Brazil

<sup>d</sup> Unité d'Antibiogénèse et Virulence Bactériennes, Université de Lyon, Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail, (ANSES), Lyon, France

<sup>e</sup> Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, Brazil

<sup>f</sup> Departamento de Doenças Dermatológicas, Infecciosas e Parasitárias, Faculdade de Medicina de São José do Rio Preto, Av. Brigadeiro Faria Lima 5416, 15090-000 São José do Rio Preto, SP, Brazil

## ARTICLE INFO

## Article history:

Received 6 February 2017

Received in revised form 23 March 2017

Accepted 6 April 2017

Available online 27 May 2017

## Keywords:

*Escherichia coli*

Extended-spectrum  $\beta$ -lactamase

InC1

Food

WGS

Brazil

## ABSTRACT

*Escherichia coli*, the main host of the CTX-M-15 extended-spectrum  $\beta$ -lactamase (ESBL) enzyme, is widely distributed and exchanged between the environment, animals and humans. Therefore, identification of *bla*<sub>CTX-M-15</sub>-positive lineages in food has a significant impact on public health. In this regard, until the end of 1990s, ESBL-producing isolates were mainly associated with hospital-acquired infections, with a predominance of SHV- and TEM-type enzymes. In recent years, a new trend has been observed among ESBL-producers, where most isolates now harbour CTX-M-type, being further isolated from community-acquired infections. Nowadays, CTX-M-15 has been recognised as the most important ESBL variant, invading virtually all human and animal compartments, leading to a global pandemic. Thus, whilst the rapid emergence and dissemination of CTX-M-15 among *E. coli* isolates has generated a large genetic reservoir from which other members of the Enterobacteriaceae family can easily acquire this resistance gene, there are an increasing number of new reservoirs and transmission mechanisms that must be investigated. In this study, we present the draft genome sequence of a CTX-M-15-producing *E. coli* ST345 isolated from commercial chicken meat in Brazil. This draft genome can be used as a reference sequence for comparative analysis among CTX-M-15-producers.

© 2017 Published by Elsevier Ltd on behalf of International Society for Chemotherapy of Infection and Cancer.

Extended-spectrum  $\beta$ -lactamases (ESBLs) remain the most important mechanism conferring resistance to broad-spectrum cephalosporins (BSCs) in *Escherichia coli*. Spread of ESBL-producing bacteria to humans through food products is a matter of concern, especially through retail meat [1]. In particular, chicken meat has recently been pointed out as a non-human source of ESBLs worldwide, whereas ceftiofur use has resulted in the selection of BSC-resistant bacteria from chickens [2]. During slaughtering, enteric bacteria from the animal's gut may contaminate the carcass and may then reach the human gut through consumption of contaminated meat or foods that come into contact with

contaminated meat or kitchen utensils [3]. Indeed, a few ESBL-producing *E. coli* isolates from human clinical samples were also identified from food-producing animals [1]. In Brazil, previous studies have reported the presence of *E. coli* producing CTX-M-type enzymes in retail chicken meat [4]. Since Brazil is the greatest chicken meat exporter in the world [5], there is an urgent need to strengthen epidemiological surveillance owing to public health concerns.

In 2014, during a surveillance study conducted to assess the occurrence of ESBL-producing bacteria in commercial chicken meat [4], a BSC-resistant CTX-M-15-producing *E. coli* isolate (Ec39764) was recovered from a sample collected at a marketplace located in São Paulo state, Southeastern Brazil.

\* Corresponding author. Fax: +55 17 3229 1777.

E-mail address: [ml.nogueira@famerp.br](mailto:ml.nogueira@famerp.br) (M.C.L. Nogueira).

The total genomic DNA of Ec39764 was sequenced using the NextSeq Platform (Illumina Inc., San Diego, CA). A total of 12 514 906 paired-end reads (150 × 150 bp) were generated with 375× coverage. In brief, the sequence reads were initially assembled de novo using Velvet v.1.2.10 (<https://www.ebi.ac.uk/~zerbino/velvet/>), and contigs were curated using Geneious software v.R10 (Biomatters Ltd., Auckland, New Zealand). Finally, the draft genome sequence was automatically annotated using PROKKA (<http://www.vicbioinformatics.com/software.prokka.shtml>).

In total, 5028 coding sequences, 288 pseudogenes, 58 tRNAs and 1 rRNA (5S) were identified in Ec39764, with a G + C content of 50.4%. Multilocus sequence typing (MLST), serotypes, plasmid replicons, acquired antimicrobial resistance genes and *E. coli* virulence genes were identified using the services MLST 1.8, SeroTypeFinder 1.1, PlasmidFinder 1.3, ResFinder 2.1 and VirulenceFinder 1.5, respectively, available from the Center for Genomic Epidemiology (<http://genomicepidemiology.org/>).

*E. coli* isolate Ec39764 belongs to ST345, and resistome analysis revealed the presence of aminoglycoside resistance genes [*aph(3')*-*Ia*, *aadA1*, *aadA2*, *strA* and *strB*], β-lactam resistance genes (*bla*<sub>TEM-18</sub> and *bla*<sub>CTX-M-15</sub>), the fluoroquinolone resistance gene *qnrS1*, the phenicol resistance gene *cmlA1*, sulphonamide resistance genes (*sul2* and *sul3*), the tetracycline resistance gene *tet(A)* and the trimethoprim resistance gene *dhfrA12*. Plasmids of incompatibility (Inc) groups X1, FIB, I1, FII and Q1 were identified, and the *bla*<sub>CTX-M-15</sub> gene was harboured by an IncX1 plasmid since both are present in the same contig. Finally, Ec39764 carried *iroN*, *cma*, *lpfA*, *iss*, *astA* and *gad* virulence genes.

In summary, here we report the first draft genome sequence of a CTX-M-15-producing *E. coli* ST345 isolate from commercial chicken meat, which can be used as a reference sequence for comparative analysis among CTX-M-15-positive lineages of *E. coli* isolated from human, animal and environmental samples.

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession MUFZ00000000. The version described in this paper is version MUFZ01000000.

## Funding

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) [2014/17184-0] and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). NL is research fellow of Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). LTC is the recipient of a postgraduate fellowship from FAPESP [2015/21325-0].

## Competing interests

None declared.

## Ethical approval

Not required.

## References

- [1] Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, et al. Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. *Clin Microbiol Infect* 2011;17:873–80.
- [2] Dutil L, Irwin R, Finley R, Ng LK, Avery B, Boerlin P, et al. Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. *Emerg Infect Dis* 2010;16:48–54.
- [3] Calbo E, Freixas N, Xercavins M, Riera M, Nicolás C, Monistrol O, et al. Foodborne nosocomial outbreak of SHV1 and CTX-M-15-producing *Klebsiella pneumoniae*: epidemiology and control. *Clin Infect Dis* 2011;52:743–9.
- [4] Casella T, Rodríguez MM, Takahashi JT, Ghiglione B, Dropa M, Assunção E, et al. Detection of *bla*<sub>CTX-M</sub>-type genes in complex class 1 integrons carried by Enterobacteriaceae isolated from retail chicken meat in Brazil. *Int J Food Microbiol* 2015;197:88–91.
- [5] United States International Trade Commission (USITC). Brazil: competitive factors in Brazil affecting U.S. and Brazilian agricultural sales in selected third country markets. Washington, DC: USITC; 2012.